

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/342666026>

Quantitative assessment and spatial distribution of avian fauna of Okhla Bird Sanctuary of Delhi NCR

Article · January 2020

CITATION

1

READS

730

2 authors, including:



[Kiranmay Sarma](#)

Guru Gobind Singh Indraprastha University

95 PUBLICATIONS 1,107 CITATIONS

SEE PROFILE



Quantitative Assessment and Spatial Distribution of Avian Fauna of Okhla Bird Sanctuary of Delhi NCR

H.S. Sarma and Kiranmay Sarma*

¹University School of Environment Management,

Guru Gobind Singh Indraprastha University, Sector 16C, Dwarka, New Delhi-110078, India

*Email: kiranmay@ipu.ac.in

Article history:

Received 30 April 2019

Received in revised form

10 January 2020

Accepted 10 January 2020

Available online

11 January 2020

Keywords:

Okhla Bird Sanctuary;

Vulnerable species;

Threatened species;

Point Count method;

Relative diversity index,

Geospatial technology

Abstract

The present study was conducted to quantitatively assess the avian fauna of Okhla Bird Sanctuary (OBS) of Delhi NCR and their spatial distribution. Point count method was applied for the study from selected 25 locations covering the entire OBS. Total number of 22 species of 10 families in 8 orders were recorded during the study, out of which 10 are migratory in nature. Two species each have been listed as vulnerable (*Aythya ferina* and *Clanga clanga*) while one species was found as near threatened (*Mycteria leucocephala*) from the total recorded species. The fully water dependant species counted was 15, partly water dependant 6 and one species was terrestrial origin. The maximum relative diversity of families was found for Anatidae (22.74) followed by Phalacrocoracidae (13.63), and Ciconiidae (9.09). The spatial distribution of total species was depicted using geospatial technology where diverse distribution was found mostly in the areas bordering Uttar Pradesh state. The area touching the NCT of Delhi side had low diverse distribution of the species.

Introduction

Wetlands are unique productive ecosystem comprising elements of both terrestrial and aquatic systems where the water table is usually at or near the surface or the land is covered by shallow water. Wetlands should support few important attributes like: (a) hydrophytes should be the predominant vegetation; (b) the soil should be hydric in nature; and (c) the underlying layer should be saturated with water or covered by shallow water at least during the growing season of each year (Cowardin *et al.*, 1979). According to Ramsar Convention (1971) wetlands shall include marsh, fen, peatland which may natural or artificial, with water may be flowing or stagnant in nature. Moreover, water can be fresh or salty including marine water bodies where depth of low tide should not be higher than 6 meter in height. Wetlands support diverse range of flora and fauna and waterbirds are one of them. Waterbirds are defined as species of birds that are

ecologically dependent on wetlands. These birds are considered to be an important health indicator of wetlands and are key parts of wetland ecosystem. Their presence, numbers and trends at a site can tell a lot about the health and quality of a wetland. Waterbirds have an important social function, providing food, recreation and tourism opportunities. Wetlands provide habitat, foods and also breeding ground for these birds (Wetland International, 2018). Waterbirds help in pest control, pollination, seed dispersal and good indicator of climate change due to their sensitive nature and thus rightly termed as bio-indicator (Harisha and Hosetti, 2009).

The study relating birds, its ecological role and need for conservation and its associated habitats, have been extensively studied all over the globe (Sidney, 1996, Tryjanowski *et al.*, 2005, Robinson, *et al.*, 2010, Figuerola and Green, 2002, Wiacek *et al.*, 2015, Rappole and Hubalek, 2003). In India, Bhat *et al.* (2009), Dar and Dar (2009), Rajashekara and Venkatesha (2010), Kumar and Gupta (2009), Thakur *et al.* (2010), Gupta *et al.* (2010), Datta (2011), Chopra and Sharma (2012), Prasad *et al.*, (2012), Gupta *et al.* (2012), Kumar and Gupta (2013), Singh *et al.* (2013), Garg *et al.* (2013), Puri and Virani (2016) have made significant contribution towards studying avian fauna migration, its conservation along with habitat and its associate wetland conservation. In the context of Okhla Bird Sanctuary significant studies have been made by Urfi (2003), Manral *et al.* (2012), Mukherjee and Sarma (2014) and Mazumdar (2017). Very few studies related to the aspects of birds have been done using geospatial technology (Swatantran *et al.*, 2012, Somveille *et al.*, 2013, Bouten *et al.*, 2013). In India Chopra *et al.* (2001) and Yadav *et al.* (2013) have made significant contributions in studying bird migration pattern with the help of geospatial tools.

Okhla Bird Sanctuary (OBS) is known as the home of migratory birds located within the territory of the Indian national capital, Delhi. The sanctuary is providing excellent contribution in terms of groundwater recharge for the city dwellers. It is also a rich source of fresh water storage for Delhi region, helps in flood control, provides groundwater recharge and livelihood for fishing community of the nearby villages. Besides the diverse aquatic plants and animals, the OBS is the home of many terrestrial plants (*Leucaena leucocephala*, *Ficus benghalensis* and *Prosopis juliflora*) and animals (*Canis aureus*, *Boselaphus tragocamelus*, and *Funambulus palmarum*), located in the heart of the city. The most attractive feature of OBS is the presence of huge flocks of beautiful winter birds which gather in huge number for resting during their non-breeding periods. These colourful birds attract a lots of nature lover during the winter season. OBS gives shelters to 302 species of birds, out of which 3 species are listed as critically endangered (*Gyps bengalensis*, *Gyps indicus* and *Aythya baeri*), 8 species are vulnerable (*Anas formosa*, *Grus antigone*, *Vanellus gregarious*, *Rynchops albicollis*, *Haliaeetus leucoryphus*, *Leptoptilos javanicus*, *Chaetornis striata* and *Ploceus megarhynchus*) and 7 species are found to be near threatened (Urfi, 2003). The Okhla Bird Sanctuary is under threat from different anthropogenic activities. An attempt has been made in the present

study to do a quantitative assessment and spatial distribution of avian fauna within Okhla Bird Sanctuary.

Materials and Methods

Study Area: Okhla Bird Sanctuary is located between 28°32'43.5"N and 28°32'56.3"N latitudes and 77°18'41.7"E and 77°18'56.6"E longitudes with altitude of 200 m above mean sea level (Mukherjee and Sarma, 2014) (Figure 1). The sanctuary falls in the districts of South West of Delhi state and Gautam Buddh Nagar district of Uttar Pradesh state. After the construction of Okhla barrage in 1987, Government of Uttar Pradesh declared it as protected area in 1990, under Wildlife protection act of 1972 (Urfi, 2003).

The study for quantifying the avian fauna of Okhla Bird Sanctuary was conducted in the month of February, 2019. Point count method (Bibby *et al.*, 2000) has been adopted for assessing the avian fauna diversity taking 25 locations covering the entire bird sanctuary. For species identification Ali (1943) has been extensively used while for assessing different status of birds, the works of Kumar *et al.*, (2013) and Mazumdar (2017) have been utilised. For calculating the relative diversity (RDi) of family the method given by Kohli (2014) is adopted.

$$RDi = \frac{\text{Number of bird species in a family}}{\text{Total number of species}} \times 100$$

For preparing the bird distribution map GCPs (ground control point) of 25 locations have been inserted in the OBS map and are linked to the field data. The total birds count in each point has been interpolated using spatial interpolation technique (kriging) in ArcGIS environment to prepare the bird distribution map.

Results and Discussion

During the present study a total number of 22 species of 10 families and 8 orders were recorded out of which 10 are migratory in nature. Two species each have been listed as vulnerable (*Aythya ferina* and *Clanga clanga*) and one near threatened (*Mycteria leucocephala*) from the total recorded species. The fully water dependant species counted was 15, partly water dependant 6 and one species was terrestrial origin (Table 1). The maximum relative diversity of families was found for Anatidae (22.74) followed by Phalacrocoracidae (13.63) and Ciconiidae (9.09) (Table 2). The spatial distribution of total species depicted diverse distribution pattern within OBS. Dense distributions of bird are found in the northeast, east and southeast parts of the sanctuary that border the Uttar Pradesh state. These sides of the bird sanctuary are comparatively less affected by the anthropogenic activities. The portion touching the Delhi state had low distribution of the

bird species due to various developmental activities taking place in this area (Figure 2). It was found from the analysis of bird distribution that about 3 percent (0.13 sq.km) of OBS area enjoys birds more than 130 numbers, located in the extreme northeast corner of the sanctuary where there are minimum human disturbances. Adjacent to this area where number of birds ranges from 116 to 130 occupies about 4 percent (0.16 sq.km) area. In the western part of the sanctuary bordering Delhi state where bird distribution was found to be scattered (below 25 species) covers more than 27 percent (1.14 sq.km) of the total sanctuary area (Table 3).




Figure 1 Location of the Okhla Bird Sanctuary




Figure 2 Distribution of avian fauna within Okhla Bird Sanctuary

Anatidae, Phalacrocoracidae and Rallidae to be found most diverse avian family in OBS. This result is the evident from other studies in wetland habitats (Urfi, 2003; Mazumdar, 2017; Vijayan *et al.*, 2006; Vyas *et al.*, 2009; Zakaria *et al.*, 2009; Ali *et al.*, 2011; Datta, 2011; Dal and Vaghela 2015; Tulasi *et al.*, 2016). The most abundant trees of the sanctuary i.e., *Leucaena leucocephala*, *Ficus benghalensis* and *prosopis juliflora* where resident birds like Cormorants and Herons use to rest for nesting purpose, and the grasses like *Typha angustifolia*, *T. Elephantina*, *Arundo donax* for feeding and for shelter (Mukherjee and Sarma, 2014). Similar results were depicted by Manral, *et al.* (2012) to support the results of this study. The course of the river Yamuna in Delhi sector is facing tremendous anthropogenic pressure leading to deteriorating water quality and changing

water regime (Rawat *et al.*, 2003; Aleem and Malik, 2005; Sengupta 2006; Trisal *et al.*, 2008). The area of OBS bordering Delhi state has been severely affected due to various development activities and other anthropogenic pressure (Manral *et al.*, 2012). Such excessive anthropogenic pressure on the riparian habitat and high load of pollutants make this stretch of Yamuna one of the most threatened riverine habitats of the world (Kumar, 2001). Observations of these factors support the results derived in the present study.

Table 1: The bird species observed in Okhla Bird Sanctuary with their respective taxonomic positions, dispersal and IUCN status and habitat

Common name	Scientific name	Family	Residential status	IUCN status	Habitat
Eurasian Marsh Harrier	<i>Circus aeruginosus</i>	Accipitridae	M	LC	W
Greater Spotted eagle	<i>Clanga clanga</i>	Accipitridae	M	VU	T
Northern shoveler	<i>Spatula clypeata</i>	Anatidae	M	LC	W
Gadwall	<i>Mareca strepera</i>	Anatidae	M	LC	W
Bar-headed goose	<i>Anser indicus</i>	Anatidae	M	LC	W
Greylag goose	<i>Anser anser</i>	Anatidae	M	LC	W
Common Pochard	<i>Aythya ferina</i>	Anatidae	M	VU	W
Indian pond heron	<i>Ardeola grayii</i>	Ardeidae	R	LC	WA
Cattle egret	<i>Bubulcus ibis</i>	Ardeidae	R	LC	WA
Great white egret	<i>Ardea alba</i>	Ardeidae	R	LC	WA
Black stork	<i>Ciconia nigra</i>	Ciconiidae	M	LC	WA
Painted stork	<i>Mycteria leucocephala</i>	Ciconiidae	R	NT	WA
Great white pelican	<i>Pelecanus onocrotalus</i>	Pelecanidae	M	LC	W
Great cormorant	<i>Phalacrocorax carbo</i>	Phalacrocoracidae	R	LC	W
Little cormorant	<i>Microcarbo niger</i>	Phalacrocoracidae	R	LC	W
Indian cormorant	<i>Phalacrocorax fuscicollis</i>	Phalacrocoracidae	R	LC	W
Greater flamingo	<i>Phoenicopterus roseus</i>	Phoenicopteridae	M	LC	W
Little grebe	<i>Tachybaptus ruficollis</i>	Podicipedidae	R	LC	W
Purple swamphen	<i>Porphyrio porphyrio</i>	Rallidae	R	LC	W
Common Coot	<i>Fulica atra</i>	Rallidae	R	LC	W
Eurasian Spoonbill	<i>Platalea leucorodia</i>	Threskiornithidae	R	LC	W
Black ibis	<i>Pseudibis papillosa</i>	Threskiornithidae	R	LC	WA

M: Migratory, R: Resident, LC: Least concern, NT: Near threaten, VU: Vulnerable, W: Water dependent, WA: Water associate, T: Terrestrial

Table 2: Relative diversity index (RDi) of avian family of Okhla Bird Sanctuary

Sl. No.	Family	Rdi	Sl. No.	Family	Rdi
1.	Anatidae	22.72	6.	Threskiornithidae	9.09
2.	Phalacrocoracidae	13.63	7.	Phoenicopteridae	4.54
3.	Rallidae	13.63	8.	Pelecanidae	4.54
4.	Ardeidae	13.63	9.	Podicipedidae	4.54
5.	Ciconiidae	9.09	10.	Accipitridae	4.54

Table 3: Spatial distribution of avian fauna in Okhla Bird Sanctuary

Sl. No.	Number of species	Area (Sq.km)	Percentage
1.	Above 130	0.13	3.15
2.	116 to 130	0.16	4.12
3.	101 to 115	0.24	6.05
4.	86 to 100	0.30	7.51
5.	71 to 85	0.45	11.14
6.	56 to 70	0.71	17.67
7.	41 to 55	0.48	12.11
8.	25 to 40	0.41	10.41
9.	Below 25	1.11	27.85
	Total	4.00	100.00

Conclusions

Okhla Bird Sanctuary is one of the few remaining wetlands of Delhi NCR. This wetland plays a vital role in providing shelter to avian fauna, attracts huge flock of migratory birds and also provides irrigation water via Agra canal to the neighbouring state of Uttar Pradesh. The OBS is under tremendous anthropogenic pressure like, urbanization, land encroachment, degradation, water and noise pollutions and so on. Noise pollution created by heavy vehicular movements, high rise buildings and electric poles and wires above OBS is affecting the avian ecosystem of the sanctuary. Moreover, Delhi side of the sanctuary is not covered under the Protected Area Network and as a result, most of the disturbing activities are prevailing in these parts. For effective conservation of avian fauna along with its associated ecosystem there should be proper management and protection in all sides of the sanctuary.

Authors Contribution: H.S. Harma (M.Sc. NRM) performed the research work, data collection and writing of the manuscript; Kiranmay Sarma (Associate Professor) has contributed in GIS part and its interpretation including the editing of manuscript and also the corresponding author.

References

- Aleem, A., Malik, A., 2005. Genotoxicity of the Yamuna river water at Okhla (Delhi), India. *Ecotoxicology and Environmental Safety*, 61(3), 404–412.
- Ali, S., 1943. *The book of Indian birds*. The Bombay Natural History Society, Bombay.
- Ali, Z., Bibi, F., Shelly, S.Y., Qazi, A., Khan, A.M., 2011. Comparative avian faunal diversity of Jiواني coastal wetlands and Taunsa barrage Wildlife Sanctuary, Pakistan. *The Journal of Animal and Plant Sciences*, 21(2), 381–387.
- Bhat, P.I., Cristopher, S.S., Hosetti, B.B., 2009. Avifaunal diversity of Anekere wetland, Karkala, Udupi district, Karnataka, India. *Journal of Environmental Biology*, 30(6), 1059-1062.
- Bibby, C.J., Burgess, N.D., Hill, D.A., Mustoe, S., 2000. *Bird census techniques*. Elsevier.
- Bouten, W., Baaij, E.W., Shamoun-Baranes, J., Camphuysen, K.C., 2013. A flexible GPS tracking system for studying bird behaviour at multiple scales. *Journal of Ornithology*, 154(2), 571-580.
- Chopra, G., Sharma, S.K., 2012. Avian biodiversity in and around major wetlands of “Lower Shivalik foothills” (India). *Nature and Science*, 10(7), 86-93.

- Chopra, R., Verma, V.K., Sharma, P.K., 2001. Mapping, monitoring and conservation of Harike wetland ecosystem, Punjab, India, through remote sensing. *International Journal of Remote Sensing*, 22(1), 89-98.
- Cowardin, L.M., Carter V., Golet C.F., LaRoe T.E., 1979. Classification of wetlands and deepwater habitat of the United States. *U.S. Fish and Wildlife Service*, Northern Prairie Wildlife Research Center.
- Dal, P., Vaghela, A.K., 2015. Preliminary survey of avifaunal diversity around Shetrunji River, Dhari, India. *Journal of Biology and Earth Sciences*, 5(1), 19-24.
- Dar. I.A., Dar. M.A., 2009. Seasonal variations of avifauna of Shallabug wetland, Kashmir. *Journal of Wetlands Ecology*, 2, 20-34.
- Datta, T., 2011. Human interference and avifaunal diversity of two wetlands of Jalpaiguri, West Bengal, India. *Journal of Threatened Taxa*, 3(12), 2253-2262.
- Figuerola, J., Green, A.J., 2002. Dispersal of aquatic organisms by waterbirds: a review of past research and priorities for future studies. *Freshwater Biology*, 47, 483-494.
- Garg, R.K., Rao, R.J., Saksena, D.N., 2013. Spatial relations of migratory birds and water quality management of Ramsagar reservoir, Datia, Madhya Pradesh, India. *Journal of Ecology and the Natural Environment*, 5(10), 335-339.
- Gupta, R., Kumar, C., Kaushik, T.K., Kumar. S., 2010. Evaluation of the extent of wetland birds in district Kaithal, Haryana, India. *Journal of Applied and Natural Science*, 2(1), 77-84.
- Gupta, R.C., Parashar, M., Kaushik, T.K., 2012. Documentation of avian diversity of Khaparwas Bird Sanctuary in Jhajjar district in Haryana, India. *International Journal of Life Sciences*, 6(1), 20-30
- Harisha, M.N., Hosetti, B.B., 2009. Diversity and distribution of avifauna of Lakkavalli range forest, Bhadra Wildlife Sanctuary, Western Ghat, India. *Ecoprint*, 16, 21-27.
- Kohli, V.K., 2014. Diversity and status of avifauna in Todgarh-Raoli Wildlife Sanctuary, Rajasthan, India. *Journal of Asia-Pacific Biodiversity*, 7(4), 401-407.
- Kumar, P., 2001. Cost of wetland conversion: A case study of floodplain wetland ecosystems along the Yamuna river corridors of Delhi region. In Report of open meeting of the Global Environmental Change Research Community, 6th-8th October 2001, Rio de Janeiro, Brazil.
- Kumar, P., Gupta, S.K., 2013. Status of wetland birds of Chhilchhila Wildlife Sanctuary, Haryana, India. *Journal of Threatened Taxa*, 5(5), 3969-3976.
- Kumar, P., Gupta, S.K., 2009. Diversity and abundance of wetland birds around Kurukshetra, India. *Our Nature*, 7, 187-192.
- Manral, U., Raha, A., Solanki, R., Hussain, S.A., Mohan, D., Talukdar, G., Veeraswami, G.G., 2012. Hydrological characteristics and flood plain vegetation of human impacted wetlands: A case study from Okhla Bird Sanctuary, National Capital Region, India. *Asian journal of conservation Biology*, 1(2), 110-119.
- Mazumdar, S., 2017. Composition of avian communities in a human-modified wetland Okhla Bird Sanctuary, India: with notes on conservation initiatives. *Proc Zool Soc.*, 1-15.
- Mukherjee, A., Sarma, K., 2014. Community structure of plant species in Okhla Bird Sanctuary, Delhi, India. *International Journal of Conservation Science*, 5(3), 397-408.
- Prasad, P.R.C., Kumari. A., Mathew. M., Thomas. R., 2012. Human bird conflicts and management issues: a case study of birds at Uppalapadu lake, Andhra Pradesh, India., *International Journal of Zoology*, Article ID 848757. p. 9.
- Puri, S.D., Virani, R.S., 2016. Diversity and status of avifauna from Bodalkasa lake in Gondia district, Maharashtra, India. *Int. J. of Life Sciences*, 4 (2), 256-262.
- Rajashekara. S., Venkatesha. M.G., 2011. Community composition of aquatic birds in lakes of Bangalore, India. *Journal of Environmental Biology*, 32 (1), 77-83.
- Ramsar handbooks., 2016. 5th edition.
- Rappole. J.H., Huba'lek, Z., 2003. Migratory birds and West Nile virus, *Journal of Applied Microbiology*, 94, 47S-58S.
- Rawat, M., Moturi, M.C.Z., Subramanian, V., 2003. Inventory compilation and distribution of heavy metals in wastewater from small-scale industrial areas of Delhi, India. *Journal of Environmental Monitoring*, 5(6), 906-912.

- Robinson, W.D., Bowlin, M., Bisson, I., Barnes, J.S., Thorup, K., Diehl, R.H., Kunz, T.H., Mabey, S., Winkler, D.W., 2010. Integrating concepts and technologies to advance the study of bird migration. *Frontiers in Ecology and the Environment*, 8(7), 354-361.
- Sengupta, B., 2006. Water quality status of Yamuna River (1999–2005), assessment and development of river basin series: ADSORBS/41/2006–07. Delhi: Central Pollution Control Board.
- Sidney A.G., 1996. Historical perspectives bird migration: methodologies and major research trajectories. *The Condor*, 98(2), 442-453.
- Singh, S., Sreeja S.N., Gupta, A.K., 2013. Ecosystem services for disaster risk reduction: a case study of wetland in East Delhi region, India. *Global Journal of Human Social Science Geography, Geo-Sciences, Environmental Disaster Management*, 13(4), 1.
- Somveille, M., Manica, A., Butchart, S.H., Rodrigues, A.S., 2013. Mapping global diversity patterns for migratory birds. *PloS one*, 8(8), e70907.
- Swatantran, A., Dubayah, R., Goetz, S., Hofton, M., Betts, M.G., Sun, M., Simard, M., Holmes, R., 2012. Mapping migratory bird prevalence using remote sensing data fusion. *Plos one*, 7(1), e28922.
- Thakur, M.L., Mattu, V.K., Lal, V.N., Sharma, H., Raj, H., Thakur, V., 2010. Avifauna of Arki Hills, Solan (Himachal Pradesh), India. *Indian Birds*, 5(6).
- Trisal, C., Tabassum, T., Kumar, R., 2008. Water quality of the river Yamuna in the Delhi stretch: Key determinants and management issues. *Clean—Soil, Air, Water*, 36(3), 306–314.
- Tryjanowski, P., Kuz'niak, S., Sparks, T.H., 2005. What affects the magnitude of change in first arrival dates of migrant birds? *J Ornithol*, 46, 200–205.
- Tulasi, P.T., Lakshmi, B.B., Laxmikanth, P., Srinivas, P., 2016. Biodiversity conservation of Tatipudi reservoir with reference to avifauna. *International Journal of Plant, Animal and Environmental Sciences*, 7(1), 45–55.
- Urfi, A.J., 2003. The birds of Okhla barrage bird sanctuary, Delhi, India, *Forktail*, 19,39-50.
- Vijayan, L., Prasad, S.N., Sridharan, N., Guptha, M.B., 2006. *Status of wetlands and wetland birds in selected districts of Tamilnadu*. Salim Ali Centre for Ornithology and Natural History, Coimbatore.
- Vyas, V., Saxena, S., Balapure, S., Shrivastava, P., 2009. *Biodiversity conservation in wetlands of Barna reservoir with reference to migratory birds*. In Proceedings of the 13th world lake conference, Wuhan: International Lake Environment Committee.
- Wetland International., 2018.
- Wiaceka, J., Polak, M., Kucharczyka, M., Bohatkiewicz, J., 2015. The influence of road traffic on birds during autumn period: implications for planning and management of road network. *Landscape and Urban Planning*, 134, 76–82.
- Yadav, M.K., Yadav, S.K., 2013. GIS applications in habitat conservation for migratory birds in Uttar Pradesh. *International Journal of Environmental Science: Development and Monitoring*, 4(2), 2231-1289.
- Zakaria, M., Rajpar, M.N., Sajap, A.S., 2009. Species diversity and feeding guilds of birds in Paya Indah Wetland Reserve, Peninsular Malaysia. *International Journal of Zoological Research*, 5(3), 86–100.